

Proof Nets for Unit-free Multiplicative-Additive Linear Logic (Extended abstract)*

DOMINIC HUGHES

Stanford University

ROB VAN GLABBEEK

Stanford University

A cornerstone of the theory of proof nets for unit-free multiplicative linear logic (MLL) is the abstract representation of cut-free proofs modulo inessential commutations of rules. The only known extension to additives, based on monomial weights, fails to preserve this key feature: a host of cut-free monomial proof nets can correspond to the same cut-free proof. Thus the problem of finding a satisfactory notion of proof net for unit-free multiplicative-additive linear logic (MALL) has remained open since the inception of linear logic in 1986. We present a new definition of MALL proof net which remains faithful to the cornerstone of the MLL theory.

1 Introduction

The beautiful theory of proof nets for unit-free multiplicative linear logic (MLL) appeared alongside the introduction of linear logic [Gir87]. A proof net is an abstract representation of a proof: the translation of cut-free proofs into proof nets identifies proofs modulo inessential commutations of rules. The identifications have since been verified as canonical from a semantic perspective, with numerous full completeness results for MLL, *e.g.* [AJ94, HO93, Loa94, Tan97, BS96, DHPP99]. Furthermore, the identifications correspond to coherences of free star-autonomous categories [BCST96].

The problem of finding a satisfactory extension of the theory of proof nets to unit-free multiplicative-additive linear logic (MALL) has remained open since the inception of linear logic [Gir87]. Progress towards a solution was made by Girard [Gir96] with a notion of MALL proof net based on monomial weights. Unfortunately, monomial proof nets fail to extend the MLL theory faithfully: a single cut-free proof may correspond to a host of monomial proof nets, and there is no natural translation of cut-free proofs into monomial proof nets. Indeed, to quote Girard, monomial proof nets are “far from being absolutely satisfactory” [Gir96]. We illustrate the problems in detail in Section 4.1.

In this paper we propose a new notion of MALL proof net (Section 2) which adheres faithfully to the original MLL

theory: we provide a simple inductive translation of cut-free proofs into cut-free proof nets, yielding the sought-after abstract representations of cut-free proofs modulo inessential commutations of rules. We define a cut-free proof net on a sequent Γ as a set of linkings on Γ satisfying a geometric correctness criterion (Definition 1), and prove that a set of linkings is the translation of a proof if and only if it is a proof net (Theorem 1).

In Section 3 we extend our proof nets to include the cut rule, and present a notion of cut elimination. Our approach to cut suffers from the same problem as Girard’s monomial proof nets: in the presence of cuts, multiple proof nets may correspond to the same proof. However, from a semantic point of view (*viz.* full completeness) the provision of abstract representations of MALL proofs modulo inessential rule commutations is crucial only in the cut-free setting. Moreover, our notion of cut elimination is simply defined, strongly normalising, and yields a category of proof nets in which $\&$ and \oplus are product and coproduct.

A crisp notion of cut-free MALL proof net is fully motivated from a proof-theoretic perspective alone. However, just as MLL has blossomed through numerous fully complete semantics via cut-free MLL proof nets, hopefully the new definition of cut-free proof net presented here will lead to a similar blossoming of MALL. Since cut-free monomial proof nets for MALL are unsatisfactory for the reasons mentioned earlier (and detailed in Section 4.1), any MALL full completeness result¹ based on them (*e.g.* [AM99], and the work in progress of Blute, Hamano and Scott on hypercoherence spaces) suffers accordingly, particularly with regard to faithfulness. We anticipate that our new definition of MALL proof net will yield cleaner and more accessible MALL full completeness results.

Relationship with Girard’s monomial proof nets. The technical starting point for our definition of proof net was Girard’s definition of monomial proof net [Gir96], and indeed we employ variants of Girard’s ingenious notions of slice and jump. Each of our proof nets translates naturally into a non-monomial Girard proof net, *i.e.*, a Girard

*This paper appeared in Proceedings 18th Annual IEEE Symposium on *Logic in Computer Science* (LICS 2003), Ottawa, Canada, June 2003, except for the appendices, which contain the proofs omitted in the LICS paper. A preliminary version of some of the material in this paper was presented in a talk at the workshop *Linear Logic 2002*, Copenhagen.

¹The original motivation for this work came as part of a project by the first author, Gordon Plotkin and Vaughan Pratt aiming to extend the full completeness of Chu spaces for MLL [DHPP99] to MALL. We have since discovered that the result does not extend.

proof net without the condition demanding that weights must be monomials. Thus one of our contributions relative to [Gir96] is the successful elimination of the monomial condition. In [Gir96] Girard remarks that he had been trying to circumvent this technical limitation since 1990, and Appendix A.1.5 of [Gir96] lists three specific problems that must be solved in any attempt to eliminate the monomial condition, *i.e.*, in any attempt to define what he calls “more liberal proof-nets”, such as ours:

Weights must be monomials. However, weights of the form $p \cup q$ will naturally occur if we want to allow more superimpositions. The present state of affairs is as follows:

- (1) *in spite of years of efforts, I never succeeded in finding the right correctness criterion for these more liberal proof-nets;*
- (2) *general boolean coefficients might be delicate to represent (on the other hand, the case we consider has a natural presentation in terms of coherent spaces);*
- (3) *normalization in the full case might be messy.*

An important stepping stone towards finding the right criterion to address (1) was to first settle the open problem of whether Girard’s criterion becomes insufficient upon relaxing the monomial condition. We show that this is indeed the case: in Section 4.2 we present a non-monomial proof structure that does not correspond to any proof, yet satisfies Girard’s criterion. We address (2) by leaving weights implied, defining a proof net on a sequent Γ as a set of axiom linkings on an extension Γ^+ of Γ with complementary pairs of cut formula occurrences. Point (3) is addressed by the fact that our definition of cut elimination is sufficiently simple that confluence and strong normalisation are immediate.

The proof that our correctness criterion captures proof translations hinges on an ordering of $\&$ ’s and \wp ’s which we call *domination*. By introducing domination we avoid the use of empires [Gir87, Gir96], thereby sidestepping the problem of stability of maximal empires ([Gir96], section 1.5.3)—the main technical problem that led Girard to resort to monomials in the first place.

MALL. By MALL we mean multiplicative-additive linear logic without units [Gir87]. Formulas are built from literals (propositional variables P, Q, \dots and their negations P^\perp, Q^\perp, \dots) by the binary connectives *tensor* \otimes , *par* \wp , *with* $\&$ and *plus* \oplus . Negation $(-)^{\perp}$ extends to arbitrary formulas by de Morgan duality. For technical convenience we take sequents to be unordered, *i.e.*, a sequent is a non-empty set of formula occurrences A_1, \dots, A_n . We omit turnstiles, which are redundant since all sequents are right-sided. Sequent are proved using the following rules:

$$\frac{\frac{\frac{\frac{\overline{P, P^\perp}}{\text{ax}} \quad \frac{\overline{P, P^\perp}}{\text{ax}}}{\otimes} \quad \frac{\overline{P, P^\perp \oplus Q}}{\oplus_1} \quad \frac{\overline{P, P^\perp}}{\text{ax}}}{\oplus_1} \quad \frac{\overline{P, P^\perp}}{\otimes}}{\otimes}}{\otimes}}{\&} \quad \frac{\overline{P \otimes P, P^\perp, P^\perp}}{\otimes} \quad \frac{\overline{P \otimes P, P^\perp, P^\perp \oplus Q}}{\otimes}}{\otimes} \quad \frac{\overline{P \otimes P, P^\perp, P^\perp \oplus Q}}{\&}}{\wp}$$

Figure 1. Example of the inductive translation of a cut-free proof into a cut-free proof net.

$$\frac{\overline{P, P^\perp}}{\text{ax}} \quad \frac{\frac{\Gamma, A \quad A^\perp, \Delta}{\Gamma, \Delta} \text{cut}}{\otimes} \quad \frac{\frac{\Gamma, A \quad B, \Delta}{\Gamma, A \otimes B, \Delta}}{\otimes} \quad \frac{\Gamma, A, B}{\Gamma, A \wp B} \wp$$

$$\frac{\frac{\Gamma, A \quad \Gamma, B}{\Gamma, A \& B} \& \quad \frac{\Gamma, A}{\Gamma, A \oplus B} \oplus_1 \quad \frac{\Gamma, B}{\Gamma, A \oplus B} \oplus_2}{\wp}$$

Here, and throughout this document, P, Q, \dots range over propositional variables, A, B, \dots over formulas, and Γ, Δ, \dots over sets of formula occurrences. In eliminating the permutation rule, we assume an implicit tracking of formula occurrences above the line of a rule to formula occurrences below the line. Without loss of generality (see [Gir87]) we restrict the axiom rule to literals.

Flavour of our approach. To give a flavour of our approach, Figure 1 shows an example of the inductive translation of a cut-free proof into one of our proof nets. The concluding proof net consists of two linkings, one drawn above the sequent, the other below. Each contains two axiom links. The proof nets further up in the derivation have one or two linkings, correspondingly above and/or below the sequent. Had we switched the order of the right-hand tensor rule and the plus rule, we would have obtained exactly the same pair of linkings; thus we identify cut-free proofs modulo a commutation of rules. Two additional translations are shown in Figure 2.


Here is an example of a proof net with four linkings:

$$\begin{array}{l} \overbrace{P \& P, Q \& Q, (Q^\perp \otimes P^\perp)}^{\otimes} \otimes \overbrace{((R \wp R) \wp (R^\perp \otimes R^\perp))}^{\wp} \\ \overbrace{P \& P, Q \& Q, (Q^\perp \otimes P^\perp)}^{\otimes} \otimes \overbrace{((R \wp R) \wp (R^\perp \otimes R^\perp))}^{\wp} \\ \overbrace{P \& P, Q \& Q, (Q^\perp \otimes P^\perp)}^{\otimes} \otimes \overbrace{((R \wp R) \wp (R^\perp \otimes R^\perp))}^{\wp} \\ \overbrace{P \& P, Q \& Q, (Q^\perp \otimes P^\perp)}^{\otimes} \otimes \overbrace{((R \wp R) \wp (R^\perp \otimes R^\perp))}^{\wp} \end{array}$$

$$\begin{array}{c}
\frac{\frac{\text{ax}}{P^\perp, P}}{\oplus_1} \\
\frac{\frac{\frac{\text{ax}}{Q, Q^\perp}}{\wp} \oplus_1 \frac{\text{ax}}{R, R^\perp}}{P^\perp, P} \oplus_2 \frac{\frac{\text{ax}}{Q^\wp Q^\perp} \oplus_1 \frac{\text{ax}}{R^\wp R^\perp}}{\oplus_2} \otimes \\
\frac{\frac{\text{ax}}{P^\perp, P} \oplus_1 \frac{\frac{\text{ax}}{Q^\wp Q^\perp} \oplus_1 \frac{\text{ax}}{R^\wp R^\perp}}{\oplus_2} \otimes}{P^\perp, P \otimes ((Q^\wp Q^\perp) \oplus (R^\wp R^\perp))} \otimes \\
\frac{P^\perp, P \otimes ((Q^\wp Q^\perp) \oplus (R^\wp R^\perp)) \otimes}{P^\perp \& P^\perp, P \otimes ((Q^\wp Q^\perp) \oplus (R^\wp R^\perp))} \&
\end{array}$$

Figure 2. Deriving the proof nets used in the example of cut elimination.

(To aid presentation, we duplicated the underlying sequent.) Our proof nets can be encoded compactly as collections of axiom links labelled with predicates (‘weights’). For example, the four-linking proof net above can be represented as follows:



To distinguish the $\&$'s, we have subscripted them. Every $\&$ -assignment (assignment of ‘left’ or ‘right’ to each of $\&_p$ and $\&_q$) determines a linking as follows: delete each axiom link whose predicate does not hold, where we read p (resp. \bar{p}) as “ $\&_p$ is assigned ‘left’ (resp. ‘right’)” (and q analogously). The reader can check that taking each of the four possible $\&$ -assignments in turn produces the four original linkings.

We sketch the idea behind our approach to cut elimination with an example. Consider the proof nets derived in Figure 2. Viewing MALL formulas as objects², and a proof net on A^\perp, B as a morphism $A \rightarrow B$, the left proof net is a morphism $P \rightarrow P \oplus P$ and the right proof net is a morphism $P \oplus P \rightarrow P \otimes ((Q^\wp Q^\perp) \oplus (R^\wp R^\perp))$. Composition, yielding a morphism $P \rightarrow P \otimes ((Q^\wp Q^\perp) \oplus (R^\wp R^\perp))$, in other words, a proof net on the sequent $P^\perp, P \otimes ((Q^\wp Q^\perp) \oplus (R^\wp R^\perp))$, proceeds as follows. First, we concatenate the two sequents into a combined sequent of four formulas (omitting commas):

$$P^\perp \quad [P \oplus P] \cdots [P^\perp \& P^\perp] \quad P \otimes ((Q^\wp Q^\perp) \oplus (R^\wp R^\perp))$$

The cut formulas are annotated with square brackets, and the dotted line represents the cut. Next, we merge the two original proof nets into a proof net on the combined sequent:

$$P^\perp \quad [P \oplus P] \cdots [P^\perp \& P^\perp] \quad P \otimes ((Q^\wp Q^\perp) \oplus (R^\wp R^\perp))$$

²The example should be accessible to readers with no knowledge of category theory: focus on the underlying cut elimination.

This new proof net has two linkings, one drawn above the sequent and one drawn below.³ Each linking consists of three axiom links. Cut elimination proceeds as follows:

$$\begin{array}{c}
P^\perp \quad [P \oplus P] \cdots [P^\perp \& P^\perp] \quad P \otimes ((Q^\wp Q^\perp) \oplus (R^\wp R^\perp)) \\
\downarrow \\
P^\perp \quad [P] \cdots [P^\perp] \quad P \otimes ((Q^\wp Q^\perp) \oplus (R^\wp R^\perp)) \\
\downarrow \\
P^\perp \quad P \otimes ((Q^\wp Q^\perp) \oplus (R^\wp R^\perp))
\end{array}$$

The first step, aside from eliminating the \oplus and $\&$ to leave a cut pair of literals $[P] \cdots [P^\perp]$, retains only one of the two original linkings. The underhanging linking is deleted because it is ‘inconsistent’: it chooses opposite arguments for the cut \oplus and $\&$ (left and right, respectively). This is an instance of our general rule for additive elimination: *retain precisely the consistent linkings*, those which choose the same argument for the cut \oplus and $\&$ (in the example above, the upper linking, which chooses left for both). The second step is the usual MLL elimination of a cut pair of literals, which we include in order to frame the example in a familiar context. Note that the end result really is a cut-free proof net, *i.e.*, the translation of a cut-free proof: its witness is a subproof of the right-hand proof of Figure 2 (the left branch of the final $\&$ -rule).

2 Cut-free MALL proof nets

In this section we introduce our definition of cut-free MALL proof net. As an *aide m emoire* we provide a summary of the definition in the box ahead. We treat cut in Section 3.

An *additive resolution* of a MALL sequent Γ is any result of deleting one argument subtree of every additive connective ($\&$ or \oplus) of Γ . Thus every remaining $\&$ and \oplus is

³The link of the original proof net on $P^\perp, P \oplus P$ appears in both the upper linking and the lower linking of the new proof net; it is duplicated in the merging process, to match the fact that the other proof net had two linkings.

$$\begin{array}{c}
\overline{\{\{P, P^\perp\}\}} \triangleright P, P^\perp \text{ ax} \\
\frac{\theta \triangleright \Gamma, A \quad \theta' \triangleright B, \Delta}{\{\lambda \cup \lambda' : \lambda \in \theta, \lambda' \in \theta'\} \triangleright \Gamma, A \otimes B, \Delta} \otimes \\
\frac{\theta \triangleright \Gamma, A, B}{\theta \triangleright \Gamma, A \wp B} \wp \quad \frac{\theta \triangleright \Gamma, A \quad \theta' \triangleright \Gamma, B}{\theta \cup \theta' \triangleright \Gamma, A \& B} \& \\
\frac{\theta \triangleright \Gamma, A}{\theta \triangleright \Gamma, A \oplus B} \oplus_1 \quad \frac{\theta \triangleright \Gamma, B}{\theta \triangleright \Gamma, A \oplus B} \oplus_2
\end{array}$$

Figure 3. The inductive translation of cut-free MALL proofs into sets of linkings.

unary. For example, two of 12 possible additive resolutions of the sequent

$$P^\perp \oplus (Q \oplus P^\perp), (P \& P) \otimes (R \oplus R), (R^\perp \otimes R) \wp R^\perp$$

are

$$\begin{array}{c}
\cancel{P^\perp} \oplus (\cancel{Q} \oplus P^\perp), (\cancel{P \& P} \otimes (\cancel{R \oplus R})), (R^\perp \otimes R) \wp R^\perp \\
P^\perp \oplus (\cancel{Q} \otimes \cancel{P^\perp}), (\cancel{P \& P} \otimes (\cancel{R \oplus R})), (R^\perp \otimes R) \wp R^\perp
\end{array}$$

Let Γ^* be an additive resolution of Γ . An **axiom link** on Γ^* is a pair of complementary literal occurrences of Γ^* . A **linking** on Γ^* is a partitioning of the set of literal occurrences of Γ^* into axiom links, *i.e.*, a set of disjoint axiom links whose union contains every literal occurrence of Γ^* . For example, there are two linkings possible on the first of the two additive resolutions depicted above:

$$\begin{array}{c}
\cancel{P^\perp} \oplus (\cancel{Q} \oplus P^\perp), (\cancel{P \& P} \otimes (\cancel{R \oplus R})), (R^\perp \otimes R) \wp R^\perp \\
\cancel{P^\perp} \oplus (\cancel{Q} \otimes \cancel{P^\perp}), (\cancel{P \& P} \otimes (\cancel{R \oplus R})), (R^\perp \otimes R) \wp R^\perp
\end{array}$$

Every additive resolution Γ^* of Γ induces an MLL sequent, namely by collapsing its additive connectives, which are unary in Γ^* . A linking λ on Γ^* , viewed as being on the induced MLL sequent, is exactly an MLL proof structure in the standard sense [Gir87], which we call the **MLL proof structure induced by λ** . For example, the MLL proof structure induced by the first of the two linkings above is:

$$\overbrace{P^\perp}, \overbrace{P \otimes R}, \overbrace{(R^\perp \otimes R) \wp R^\perp}$$

A **linking on a MALL sequent Γ** is a linking on an additive resolution of Γ . Write $\Gamma \upharpoonright \lambda$ for the additive resolution associated with a linking λ . Every cut-free MALL proof of Γ defines a set of linkings on Γ by a simple induction, as in Figure 3, where $\theta \triangleright \Gamma$ is the judgement “ θ is a set of linkings on Γ ”. (We use the implicit tracking of literal occurrences downwards through rules.) The base case ax is a singleton set of linkings whose only linking comprises a single axiom link, between P and P^\perp . Examples of the inductive translation of cut-free proofs into sets of linkings

were presented in Figures 1 and 2. Note that if a cut-free proof Π' can be obtained from Π by a series of rule commutations, then Π and Π' translate to the same set of linkings.

Geometric characterisation of proof translations. We present a geometric characterisation of those sets of linkings that arise as the translations of cut-free MALL proofs, and call them *proof nets*. Analogous to [Gir96], as a stepping stone to the definition of a proof net, we introduce the less restrictive notion of a *proof structure*.

A **&-resolution Γ^*** of a sequent Γ is any result of deleting one argument subtree of every $\&$ of Γ . A linking λ on Γ is **on Γ^*** if every literal occurrence of λ is in Γ^* . A set of linkings θ on Γ is a **proof structure** on Γ if it satisfies

(P1) *For every &-resolution Γ^* of Γ , exactly one linking of θ is on Γ^* .*⁴

We invite the reader to verify (P1) for the sets of linkings in Figures 1 and 2. Any proof structure can be represented compactly as a set of axiom links labelled with predicates (‘weights’), using the encoding described on the third page of the Introduction. In Section 4 we relate our proof structures to those of Girard.

The second requirement for a set of linkings θ to be a proof net is “pointwise MLL correctness”:

(P2) *Every linking of θ induces an MLL proof net.*

In other words, for each linking $\lambda \in \theta$, the MLL proof structure induced by λ is an MLL proof net, in the usual sense [Gir87, DR89]. To be self-contained, we characterise (P2) explicitly below.

Henceforth view a sequent Γ as a graph: a disjoint union of parse trees, with literals above. For a linking λ on Γ obtain the **graph \mathcal{G}_λ of λ** from the additive resolution $\Gamma \upharpoonright \lambda$ (a subgraph of Γ) by adding each axiom link a of λ as a vertex above $\Gamma \upharpoonright \lambda$, with edges from a down to its two literal occurrences. A **switching** of a linking λ on Γ is any subgraph of \mathcal{G}_λ obtained by deleting one of the two argument edges of each \wp -vertex. (P2) holds if and only if every switching of every linking of θ is a tree (acyclic and connected).

We require some auxiliary concepts to state our third and last requirement for a set of linkings to be a proof net. A set of linkings Λ **toggles** a $\&$ -occurrence w of Γ if both arguments of w are present in $\bigcup_{\lambda \in \Lambda} \Gamma \upharpoonright \lambda$. An axiom link a **depends on w within Λ** if, within Λ , a can be made to vanish by toggling w alone: there are $\lambda, \lambda' \in \Lambda$ such that $a \in \lambda$, $a \notin \lambda'$, and w is the only $\&$ toggled by $\{\lambda, \lambda'\}$.

⁴Therefore, a proof structure on Γ is a maximal clique in the coherence space of linkings on Γ with incoherence $\lambda \succ \lambda'$ iff there exists a $\&$ -resolution Γ^* of Γ such that both λ and λ' are on Γ^* .

Additive resolution of Γ : any result of deleting one argument subtree of every $\&$ or \oplus of Γ . ($\&$ -**resolution** analogously.)

Axiom link: pair of complementary literal occurrences.

Linking λ on Γ : partitioning of the set of literal occurrences in an additive resolution $\Gamma \upharpoonright \lambda$ of Γ into axiom links.

Graph \mathcal{G}_λ : $\Gamma \upharpoonright \lambda + \lambda$ + edges from each axiom link in λ to its two literal occurrences in $\Gamma \upharpoonright \lambda$.

Switching of a linking λ : any subgraph of \mathcal{G}_λ obtained by deleting one of the two argument edges of each \wp -vertex.

A set of linkings Λ **toggles** a $\&$ -occurrence w of Γ if both arguments of w are present in $\bigcup_{\lambda \in \Lambda} \Gamma \upharpoonright \lambda$.

An axiom link a **depends on** w **within** Λ if $\exists \lambda, \lambda' \in \Lambda$ such that $a \in \lambda$, $a \notin \lambda'$, and w is the only $\&$ toggled by $\{\lambda, \lambda'\}$.

Graph \mathcal{G}_Λ : $\bigcup_{\lambda \in \Lambda} \mathcal{G}_\lambda$ + **jump** edges between each axiom link in Λ and any $\&$ on which it depends within Λ .

Switch edge of a $\&$ - or \wp -vertex x in \mathcal{G}_Λ : any argument or jump edge of x .

Switching cycle of Λ : a (non self-intersecting) cycle in \mathcal{G}_Λ containing at most one switch edge of each $\&$ and \wp .

A set of linkings θ is a **proof net** if it satisfies

(P1) For every $\&$ -resolution Γ^* of Γ , exactly one linking of θ is on Γ^* .

(P2) Every switching of every linking of θ is a tree (acyclic and connected).

(P3) Every set Λ of two or more linkings of θ toggles a $\&$ that is not in any switching cycle of Λ .⁶

EXAMPLE 1 Consider the two linkings

$$\begin{aligned} \lambda_1 : & \quad P^\perp \oplus (Q \oplus P^\perp), (P \& P) \otimes (R \oplus R), (R^\perp \otimes R) \wp R^\perp \\ \lambda_2 : & \quad P^\perp \oplus (Q \oplus P^\perp), (P \& P) \otimes (R \oplus R), (R^\perp \otimes R) \wp R^\perp \end{aligned}$$

Here are λ_1 and λ_2 on their respective additive resolutions:

$$\begin{aligned} \lambda_1 : & \quad \cancel{P^\perp} \oplus (\cancel{Q} \oplus \cancel{P^\perp}), (\cancel{P} \& \cancel{P}) \otimes (\cancel{R} \oplus \cancel{R}), (\cancel{R^\perp} \otimes \cancel{R}) \wp \cancel{R^\perp} \\ \lambda_2 : & \quad P^\perp \oplus (\cancel{Q} \oplus \cancel{P^\perp}), (\cancel{P} \& \cancel{P}) \otimes (\cancel{R} \oplus \cancel{R}), (\cancel{R^\perp} \otimes \cancel{R}) \wp \cancel{R^\perp} \end{aligned}$$

Let w be the $\&$ of the sequent, and let $\Lambda = \{\lambda_1, \lambda_2\}$. The axiom link between the left-most R^\perp and the left-most R depends on w within Λ : it is present in $\lambda_1 \in \Lambda$ but not in $\lambda_2 \in \Lambda$, and w is the only $\&$ toggled by $\{\lambda_1, \lambda_2\}$. The axiom link between the right-most R and R^\perp does not depend on w within Λ , since it is present in both λ_1 and λ_2 . It is the only one of the 5 axiom links in Λ (more precisely, in $\bigcup \Lambda$) that does not depend on w within Λ .

We now extend the definition of the graph of a linking to the graph of a set of linkings. Given a set Λ of linkings on Γ , obtain the **graph** \mathcal{G}_Λ of Λ from $\bigcup_{\lambda \in \Lambda} \mathcal{G}_\lambda$ by adding, for every $\&$ -vertex w and every axiom link a depending on w within Λ , an edge between w and a . Each edge of the latter form, between a $\&$ -vertex w and an axiom link, is called a **jump** of w . Figure 4 shows $\mathcal{G}_{\{\lambda_1, \lambda_2\}}$ for λ_1 and λ_2 of Example 1, with four jumps (the curved edges). In drawing an axiom link \sqcap , we view the horizontal section as a vertex, and the two verticals as edges. We overlap edges from axiom links coming down into the same literal occurrence (*i.e.*, \neg means \neg). There is no jump to the right-most axiom link, since it does not depend on the $\&$ within $\{\lambda_1, \lambda_2\}$. Note that if $\Lambda \subseteq \Lambda'$, then \mathcal{G}_Λ is a subgraph of $\mathcal{G}_{\Lambda'}$, and that for any linking λ , $\mathcal{G}_{\{\lambda\}}$ is precisely \mathcal{G}_λ defined on the previous page. ($\mathcal{G}_{\{\lambda\}}$ has no jumps, since no $\&$ is toggled.)




Figure 4. The graph $\mathcal{G}_{\{\lambda_1, \lambda_2\}}$ of Example 1.

A **switch edge** of a $\&$ - or \wp -vertex x of \mathcal{G}_Λ is an edge between x and one of its arguments, or a jump of x (if x is a $\&$). A **switching cycle** of a set of linkings Λ is a cycle in \mathcal{G}_Λ containing at most one switch edge of each $\&$ and \wp . (We do not permit a cycle to intersect itself.) For example, in $\mathcal{G}_{\{\lambda_1, \lambda_2\}}$ of Figure 4, the cycle “ $\& \rightarrow \otimes \rightarrow \oplus \rightarrow$ left- $R \rightarrow$ left- $\{R, R^\perp\} \xrightarrow{\text{jump}} \&$ ” contains only one switch edge of the $\&$, and is therefore a switching cycle of $\{\lambda_1, \lambda_2\}$ of Example 1.

DEFINITION 1 A set θ of linkings on a MALL sequent Γ is a **cut-free proof net** if it satisfies (P1), (P2)⁵ and:

(P3) Every set Λ of two or more linkings of θ toggles a $\&$ that is not in any switching cycle of Λ .⁶

EXAMPLE 2 The set of linkings $\{\lambda_1, \lambda_2\}$ of Example 1 is not a proof net. It fails (P3) since $\mathcal{G}_{\{\lambda_1, \lambda_2\}}$ (Figure 4) contains a switching cycle through the $\&$.

⁵By dropping connectedness from (P2) we obtain a cut-free proof net for MALL with the MIX rule (hypotheses Γ and Δ , conclusion Γ, Δ).

⁶In fact, one need only verify (P3) for those Λ which are **saturated**, namely, such that any strictly larger subset of θ toggles more $\&$'s than Λ . Note that there is exactly one saturated set of linkings in θ for each **partial** $\&$ -**resolution** of Γ , the latter being any result of deleting one argument subtree of some of the $\&$'s of Γ .


$$\begin{array}{c}
\frac{}{\{\{P, P^\perp\}\} \triangleright [] P, P^\perp} \text{ax} \\
\theta \triangleright [\Sigma, \Omega] \Gamma, A \quad \theta' \triangleright [\Sigma, \Omega'] \Gamma, B \quad \& \\
\frac{}{\theta \cup \theta' \triangleright [\Sigma, \Omega, \Omega'] \Gamma, A \& B} \\
\theta \triangleright [\Omega] \Gamma, A \quad \oplus_1 \\
\frac{}{\theta \triangleright [\Omega] \Gamma, A \oplus B} \\
\theta \triangleright [\Omega] \Gamma, A \quad \oplus_2 \\
\frac{}{\theta \triangleright [\Omega] \Gamma, A \oplus B} \\
\theta \triangleright [\Omega] \Gamma, A, B \quad \wp \\
\frac{}{\theta \triangleright [\Omega] \Gamma, A \wp B}
\end{array}
\quad
\begin{array}{c}
\frac{\theta \triangleright [\Omega] \Gamma, A \quad \theta' \triangleright [\Omega'] A^\perp, \Delta}{\{\lambda \cup \lambda' : \lambda \in \theta, \lambda' \in \theta'\} \triangleright [\Omega, \Omega', A * A^\perp] \Gamma, \Delta} \text{cut} \\
\theta \triangleright [\Omega] \Gamma, A \quad \theta' \triangleright [\Omega'] B, \Delta \quad \otimes \\
\frac{}{\{\lambda \cup \lambda' : \lambda \in \theta, \lambda' \in \theta'\} \triangleright [\Omega, \Omega'] \Gamma, A \otimes B, \Delta} \\
\theta \triangleright [\Omega] \Gamma, B \quad \oplus_2 \\
\frac{}{\theta \triangleright [\Omega] \Gamma, A \oplus B} \\
\theta \triangleright [\Omega] \Gamma, A, B \quad \wp \\
\frac{}{\theta \triangleright [\Omega] \Gamma, A \wp B}
\end{array}$$

Figure 5. Rules for deriving sequentialisable sets of linkings on MALL cut sequents.

EXAMPLE 3 Consider the pair of linkings on the sequent $\Gamma \equiv P^\perp \& P^\perp, P \oplus P$ obtained as follows:

$$\frac{\frac{\frac{}{P^\perp, P} \text{ax}}{\frac{}{P^\perp, P \oplus P} \oplus_2} \quad \frac{\frac{}{P^\perp, P} \text{ax}}{\frac{}{P^\perp, P \oplus P} \oplus_1}}{\frac{}{P^\perp \& P^\perp, P \oplus P} \&}$$

Let λ_1 and λ_2 be the upper- and lower linking respectively (each having just one axiom link). We shall verify that $\{\lambda_1, \lambda_2\}$ is a cut-free proof net. Γ has two $\&$ -resolutions, $\Gamma_1^* \equiv P^\perp \& P^\perp, P \oplus P$ and $\Gamma_2^* \equiv P^\perp \& P^\perp, P \oplus P$. (P1) holds, since $\{\lambda_1, \lambda_2\}$ contains exactly one linking on Γ_i^* , namely λ_i . Here are the graphs \mathcal{G}_{λ_1} , \mathcal{G}_{λ_2} , and $\mathcal{G}_{\{\lambda_1, \lambda_2\}}$:



Each λ_i has just one switching, namely \mathcal{G}_{λ_i} . Since each \mathcal{G}_{λ_i} is a tree, (P2) holds. Finally, (P3) holds since $\{\lambda_1, \lambda_2\}$ toggles the $\&$, and the $\&$ is not in any switching cycle of $\{\lambda_1, \lambda_2\}$.

THEOREM 1 A set of linkings is the translation of a cut-free proof iff it is a cut-free proof net.

By a simple induction, the translation of a cut-free proof is a cut-free proof net. The proof of the converse reduces to a simple induction on the number of \wp 's and $\&$'s, spelled out in Appendix B, once we prove the *Separation Lemma*: for

any cut-free proof net θ , if \mathcal{G}_θ has a \wp or $\&$, then it has a \wp or $\&$ that separates. Here a \wp - or $\&$ -vertex x *separates* if it is not an argument (*i.e.*, is an outermost connective), or it is the argument of y and deleting the edge between x and y disconnects \mathcal{G}_θ . We prove the Separation Lemma via an ordering on $\&$'s and \wp 's which we call *domination*⁷, a concept reminiscent of the ordering induced by the notion of an *empire* of [Gir96], but different in an essential way. The details are in Appendix A. The proof in the case of MIX (see footnote 5) requires only minor changes.

3 Cut

A *cut* is a pair $\{A, A^\perp\}$ of complementary MALL formulas. We write $A * A^\perp$ for $\{A, A^\perp\}$, and treat $A * A^\perp$ akin to a MALL formula, referring to $*$ as the *cut connective*. (In the cut elimination example in the Introduction we drew a cut $A * A^\perp$ informally as $[A] \cdots [A^\perp]$.) A *cut sequent* is a non-empty set of occurrences of MALL formulas and cuts. A *cut-additive resolution* of a cut sequent Δ is any result of deleting some cuts from Δ and one argument subtree of every remaining additive connective ($\&$ or \oplus). Thus every remaining $\&$ and \oplus is unary. For example, here is a cut sequent followed by one of its cut-additive resolutions:

$$\begin{array}{c}
P \otimes P, Q * Q^\perp, P^\perp \oplus Q, (R \oplus S) * (R^\perp \& S^\perp) \\
P \otimes P, \cancel{Q * Q^\perp}, P^\perp \oplus \cancel{Q}, \cancel{(R \oplus S) * (R^\perp \& S^\perp)}
\end{array}$$

An *axiom link* on a cut-additive resolution Δ^* of a cut sequent Δ is a pair of complementary literal occurrences of Δ^* . A *linking* on Δ^* is a partitioning of the literal occurrences of Δ^* into axiom links, *i.e.*, a set of disjoint axiom links on Δ^* whose union contains every literal occurrence of Δ^* . A *linking on* Δ is a linking on a cut-additive resolution of Δ . We write $\Delta \upharpoonright \lambda$ for the cut-additive resolution associated with a linking λ .

Write $[\Omega] \Gamma$ for the cut sequent obtained by taking the disjoint union of a set Ω of cut occurrences and a MALL sequent Γ . A set of linkings on $[\Omega] \Gamma$ is *sequentialisable* if it can be derived from the rules in Figure 5, in which $\theta \triangleright [\Omega] \Gamma$ is the judgement “ θ is a sequentialisable set of linkings on the cut sequent $[\Omega] \Gamma$ ”. (We once again use the

⁷Unrelated to domination in flowgraphs.

$$\begin{array}{c}
\frac{\overline{P, P^\perp} \text{ ax}}{\overline{P, P^\perp} \text{ cut}} \quad \frac{\overline{P, P^\perp} \text{ ax}}{\overline{P, P^\perp} \text{ cut}}}{\overline{P, P^\perp * P, P^\perp} \text{ cut}} \quad \frac{\overline{P, P^\perp} \text{ ax}}{\overline{P, P^\perp} \text{ cut}} \quad \frac{\overline{P, P^\perp} \text{ ax}}{\overline{P, P^\perp} \text{ cut}}}{\overline{P, P^\perp * P, P^\perp} \text{ cut}} \\
\hline
\overline{P, P^\perp * P, P^\perp * P, P^\perp \& P^\perp} \&
\end{array}$$

$$\begin{array}{c}
\frac{\overline{P, P^\perp} \text{ ax}}{\overline{P, P^\perp} \text{ cut}} \quad \frac{\overline{P, P^\perp} \text{ ax}}{\overline{P, P^\perp} \text{ cut}}}{\overline{P, P^\perp * P, P^\perp} \text{ cut}} \quad \frac{\overline{P, P^\perp} \text{ ax}}{\overline{P, P^\perp} \text{ cut}} \quad \frac{\overline{P, P^\perp} \text{ ax}}{\overline{P, P^\perp} \text{ cut}}}{\overline{P, P^\perp * P, P^\perp} \text{ cut}} \\
\hline
\overline{P, P^\perp * P, P^\perp \& P^\perp} \&
\end{array}$$

Figure 6. Examples of the translation of a proof with cuts.

implicit tracking of literal occurrences downwards through rules.) The base case ax is a single linking with a single axiom link and no cuts. Figure 6 shows two examples. Each of the conclusions is a set of two linkings, one drawn above the cut sequent and one drawn below. The only difference between the derivations is the final $\&$ -rule. The left application keeps the cuts in the hypotheses separate (an instance of the $\&$ -rule taking Σ empty and $\Omega = \Omega' = P^\perp * P$), whereas the right application superimposes the two cuts ($\Sigma = P^\perp * P$ and Ω, Ω' empty).

Any derivation of a set of linkings using the rules of Figure 5 projects in an obvious way to a MALL proof, namely, by restricting to the underlying sequents (*viz.*, read Γ for $\theta \triangleright [\Omega] \Gamma$). For example, the two derivations of Figure 6 each yield the same MALL proof of $P, P^\perp \& P^\perp$.

Write $\Pi \rightsquigarrow \theta$ if Π is the MALL proof obtained from a derivation of a set of linkings θ , and say that Π is a **sequentialisation** of θ . If a MALL proof Π' can be obtained from Π by a series of rule commutations in which no $\&$ -rules are moved upwards, then Π and Π' are sequentialisations of the same set of linkings. In the cut-free case, \rightsquigarrow is a function from proofs to sets of linkings, exactly the translation defined in Figure 3. In the presence of cuts, more than one set of linkings may correspond to the same proof. For example, since the two derivations in Figure 6 have the same underlying MALL proof (say Π), the concluding sets of linkings (say θ and θ') have a common sequentialisation: $\Pi \rightsquigarrow \theta$, $\Pi \rightsquigarrow \theta'$, and $\theta \neq \theta'$.

We can of course extend the cut-free translation of proofs by always choosing Σ to be empty in the $\&$ -rule (*i.e.*, “never superimpose cuts”). However, our notion of proof net defined below, which characterises sequentialisability, does not characterise the image of this translation, since there would exist sequentialisable sets of linkings that are not

proof translations, such as

$$\overline{P \oplus P, P^\perp * P, P^\perp \& P^\perp}$$

Moreover, under this convention two proofs that differ only in a commutation of cut and $\&$ -rules would be translated to different sets of linkings.

Note that the alternative of taking Σ maximal (*i.e.*, “superimpose as many cuts as possible”) does not define a canonical function from proofs to sets of linkings, since there may be a choice of how to make the identifications. The following two $\&$ -rules illustrate such a choice.

$$\frac{\overline{P, P^\perp * P, P^\perp * P, P^\perp} \quad \overline{P, P^\perp * P, P^\perp * P, P^\perp}}{\overline{P, P^\perp * P, P^\perp * P, P^\perp \& P^\perp} \&}$$

$$\frac{\overline{P, P^\perp * P, P^\perp * P, P^\perp} \quad \overline{P, P^\perp * P, P^\perp * P, P^\perp}}{\overline{P, P^\perp * P, P^\perp \& P^\perp} \&}$$

Girard was aware of this issue in the context of monomial proof nets; see Appendix A.1.6 of [Gir96].

Geometric characterisation of sequentialisability. In the presence of cut, we update all the auxiliary definitions of Section 2 ($\&$ -resolution, \mathcal{G}_Λ , switching cycle, *etc.*) by substituting “cut sequent” for “sequent” and “cut-additive resolution” for “additive resolution” throughout.

DEFINITION 2 A set θ of linkings on a cut sequent Δ is a **proof net** if:

- (P0) At least one literal occurrence of every cut is in θ (*i.e.*, in some axiom link of some linking of θ).
- (P1) For every $\&$ -resolution Δ^* of Δ , exactly one linking of θ is on Δ^* .
- (P2) Every switching of every linking of θ is a tree (acyclic and connected⁸).
- (P3) Every set Λ of two or more linkings of θ toggles a $\&$ that is not in any switching cycle of Λ .

θ is a **proof structure** if it satisfies (P0) and (P1).

Note that (P1)–(P3) are inherited from the cut-free case.

THEOREM 2 (SEQUENTIALISATION) A set of linkings is sequentialisable iff it is a proof net.

The proof is essentially the same as the proof of Theorem 1; the cut connective $*$ is akin to an outermost \otimes .

⁸By dropping connectedness, we obtain a proof net for MALL augmented by the MIX rule.

$$\begin{array}{c}
\begin{array}{c} \text{id} \\ \vdots \\ P, Q, Q^\perp \otimes P^\perp \\ A \end{array} \quad \begin{array}{c} \text{tw} \\ \vdots \\ P, Q, Q^\perp \otimes P^\perp \\ A \end{array} \\
\hline
\begin{array}{c} P, Q, (Q^\perp \otimes P^\perp) \otimes A \\ \vdots \\ P, Q, (Q^\perp \otimes P^\perp) \otimes A \end{array} \quad t \\
\hline
\begin{array}{c} P, Q \& Q, (Q^\perp \otimes P^\perp) \otimes A \\ \vdots \\ P, Q \& Q, (Q^\perp \otimes P^\perp) \otimes A \end{array} \quad q
\end{array}
\quad
\begin{array}{c}
\begin{array}{c} \text{tw} \\ \vdots \\ P, Q, Q^\perp \otimes P^\perp \\ A \end{array} \quad \begin{array}{c} \text{tw} \\ \vdots \\ P, Q, Q^\perp \otimes P^\perp \\ A \end{array} \\
\hline
\begin{array}{c} P, Q, (Q^\perp \otimes P^\perp) \otimes A \\ \vdots \\ P, Q, (Q^\perp \otimes P^\perp) \otimes A \end{array} \quad t \\
\hline
\begin{array}{c} P, Q \& Q, (Q^\perp \otimes P^\perp) \otimes A \\ \vdots \\ P, Q \& Q, (Q^\perp \otimes P^\perp) \otimes A \end{array} \quad p
\end{array}$$

Figure 7. The proof Π_{tqp} . (We omit the unique cut-free proof of $P, Q, Q^\perp \otimes P^\perp$.)

Cut elimination. Let θ be a set of linkings on the cut sequent Δ , and let $A * A^\perp$ be a cut of Δ . Define the *elimination* of $A * A^\perp$ as follows.

- If A is a literal, delete $A * A^\perp$ from Δ , and replace any pair of axiom links $\{l, A\}, \{A^\perp, l'\}$ in a linking of θ (l and l' being other occurrences of A^\perp and A respectively) with the axiom link $\{l, l'\}$.
- If $A = A_1 \otimes A_2$ and $A^\perp = A_1^\perp \wp A_2^\perp$ (or vice versa), replace $A * A^\perp$ with two cuts $A_1 * A_1^\perp$ and $A_2 * A_2^\perp$. Retain all the original linkings.
- If $A = A_1 \& A_2$ and $A^\perp = A_1^\perp \oplus A_2^\perp$ (or vice versa) replace $A * A^\perp$ with two cuts $A_1 * A_1^\perp$ and $A_2 * A_2^\perp$. Retain precisely the ‘consistent’ linkings: delete any linkings λ such that in $\Delta \upharpoonright \lambda$ the (now unary) $\&$ and \oplus take opposite arguments (*i.e.*, such that the right argument of the $\&$ is in $\Delta \upharpoonright \lambda$ and the left argument of the \oplus is in $\Delta \upharpoonright \lambda$, or vice versa). Finally, ‘garbage collect’ by deleting $A_i * A_i^\perp$ if no literal occurrence of $A_i * A_i^\perp$ is in any of the remaining linkings.

An example of cut elimination was presented in the Introduction.

PROPOSITION 1 *Eliminating a cut from a proof net yields a proof net.*

Proposition 1 is proved in Appendix C.

THEOREM 3 *Cut elimination of proof nets is strongly normalising.*

Proof. Confluence is immediate from the definition; cut elimination reduces the size of the cut sequent, and is therefore strongly normalising. \square

A category of proof nets. Our cut elimination allows us to define a category of MALL proof nets. Objects are MALL formulas, and a morphism $A \rightarrow B$ is a cut-free proof net on the sequent A^\perp, B . The composition of $\theta : A \rightarrow B$ and $\theta' : B \rightarrow C$ is the normal form of the proof net $\{\lambda \cup \lambda' : \lambda \in \theta, \lambda' \in \theta'\}$ on $A^\perp, B * B^\perp, C$. Composition is associative, since cut elimination is strongly normalising. The identity $A \rightarrow A$ contains a linking λ on A^\perp, A iff λ matches the i^{th} literal of A^\perp with the i^{th} literal

of A . This category has the structure of a star-autonomous category minus the units; $\&$ is product, and \oplus is coproduct.


4 Girard’s monomial proof nets

In this section we relate our MALL proof nets to the monomial proof nets of Girard [Gir96].

4.1 Monomial proof nets are unsatisfactory

We give a detailed account of how monomial proof nets [Gir96] fail to provide abstract representations of cut-free MALL proofs modulo inessential commutations of rules. A single cut-free proof may correspond to a host of monomial proof nets, and there is no natural translation of cut-free MALL proofs into monomial proof nets. (The reader unfamiliar with monomial proof nets should be able to follow the general shape of the argument.)

Consider the following pair of cut-free monomial proof nets:



(Eigenvariables associated with $\&$ ’s are shown as subscripts; we omit implied weights.) These two monomial proof nets correspond to the same proof. The second monomial proof net has two forms of redundancy relative to the first: (i) the $\&$ with eigenweight q has been replaced by two similar ‘copies’, and (ii) the axiom link of weight p has been split into two.

Even if one attempts to fix a choice of representation (*e.g.* favouring the first monomial proof net above over the






Figure 8. Girard's correctness criterion is insufficient without monomials: this (abbreviated) non-monomial Girard proof net is not sequentialisable.

second), one still runs into difficulty. As a concrete illustration, we exhibit cut-free proofs Π_α and monomial proof nets θ_β for which the binary relation of sequentialisation is



Define Π_{tqp} to be the proof shown in Figure 7, where $A = (R \wp R) \wp (R^\perp \otimes R^\perp)$, id denotes the identity proof and tw denotes the twist proof. Let Π_{tpq} be the result of commuting rules q and p in Π_{tqp} , let Π_{tqp} be the result of commuting t and q in the right half of Π_{tpq} , and let Π_{ptq} be the result of commuting t and p in the right half of Π_{tpq} . Define the monomial proof nets θ_β as follows. To specify θ_β it suffices to present a configuration of weighted axiom links. On P and Q literals, fix the configuration as below-left:



We have taken as eigenweights the labels of the $\&$ -rules of the Π_α . The configuration of axiom links on A will be a disjoint union of axiom links in the identity and twist configurations: id_w and tw_w (above-right) denote a pair of axiom links of weight w in the identity and twist configurations, respectively. We specify the θ_β by the following disjoint

unions of weighted identity and twist configurations on A :

$$\begin{aligned} \theta_{\bar{p}, p\bar{q}} &: \text{id}_{p\bar{q}} \sqcup \text{tw}_{\bar{p}} \sqcup \text{tw}_{p\bar{q}} \\ \theta_{\bar{q}, p\bar{q}} &: \text{id}_{p\bar{q}} \sqcup \text{tw}_{\bar{q}} \sqcup \text{tw}_{p\bar{q}} \\ \theta_{\bar{p}\bar{q}, p\bar{q}, p\bar{q}} &: \text{id}_{p\bar{q}} \sqcup \text{tw}_{\bar{p}\bar{q}} \sqcup \text{tw}_{p\bar{q}} \sqcup \text{tw}_{p\bar{q}} \end{aligned}$$

(By redundancies of type (i) and (ii) illustrated earlier, there are of course a host of other monomial proof nets θ_β which are parodies of the three above.) Since the Π_α are equivalent modulo inessential rule commutations, any satisfactory theory of proof nets should provide a canonical representation uniting all of them. With monomial proof nets one would have to close under the sequentialisation relation between proofs and monomial proof nets depicted earlier, thereby creating a matching between the set of proofs Π_α and the set of monomial proof nets θ_β , and then artificially choose a representative from amongst the θ_β .

By contrast, in our setting we associate the same proof net with each Π_α : the four-linking proof net on the second page of the Introduction. Thus we preserve the spirit of MLL proof nets by providing an abstract representation of all of the Π_α in one.

4.2 Girard's criterion is insufficient without monomials

A key stepping stone towards our formulation of a new definition of proof net was to first settle the open problem of whether Girard's proof net correctness criterion [Gir96] becomes insufficient upon relaxing the dependency condition, which is the requirement that every weight be a monomial. The answer is yes: in Figure 8 we present a cut-free non-monomial Girard proof net θ which is not sequentialisable. By **non-monomial Girard proof net** we mean a proof net as in [Gir96] but for the omission of the dependency condition. Strictly speaking θ is merely an abbreviation of a

non-monomial Girard proof net: view each p_i as an eigenvariable and split each \oplus into a separate \oplus_1 and \oplus_2 ; formulas and remaining weights are implied.

Figure 8 also encodes one of our proof structures θ , via the notion of weight described on the third page of the Introduction. It is not a proof net, since (P3) fails: \mathcal{G}_θ contains a switching cycle passing through all four $\&$'s (follow the four jumps $\&_{p_i}$ to the axiom link $\{R_{i+1}^\perp, R_{i+1}\} \pmod{4}$).

4.3 Mapping monomial proof structures to ours

Let a *non-monomial Girard proof structure* be a proof structure as in [Gir96] but for the omission of the dependency condition. Define a non-monomial Girard proof structure to be *compact* if (a) any non-literal formula occurrence is the conclusion of exactly one link, except that a formula $A \oplus B$ may be the conclusion of both a \oplus_1 - and a \oplus_2 -link, and (b) any two literal occurrences constitute the conclusions of at most one axiom link. Each non-monomial Girard proof structure, and thus also each monomial one, can be collapsed into a compact non-monomial Girard proof structure by identifying, along with their premises, links of the same type with the same conclusion(s), and summing the weights of links and formulas so identified. This collapse does not preserve the dependency condition. Any compact non-monomial Girard proof **structure** can be obtained as the collapse of a (monomial) Girard proof **structure**.

Compact non-monomial Girard proof structures are in bijection with our proof structures. The counterpart of Girard's "technical condition" is implied by (P1) and our definition of a set of linkings in terms of additive resolutions. The ~~surjective~~ map from (monomial) Girard proof structures to our proof structures obtained by composing the collapse and the bijection preserves the property of being a sequentialisation of a particular MALL proof.

Given a set of linkings θ on a sequent Γ and a subset $\Lambda \subseteq \theta$, let $\mathcal{G}_\Lambda^\theta$ be defined as \mathcal{G}_Λ , but with jump edges between every $\&$ -vertex $w \in \mathcal{G}_\Lambda$ and every axiom link $a \in \mathcal{G}_\Lambda$ depending on w *within* θ . Note that $\mathcal{G}_\Lambda = \mathcal{G}_\Lambda^\Lambda$. Define the variant (P2*) of (P2) by using $\mathcal{G}_{\{\lambda\}}^\theta$ in place of \mathcal{G}_λ in the definition of a switching of λ , also deleting all but one switch edge of each $\&$. (P2*) clearly implies (P2), since it involves more switchings. In fact, (P2*) is strictly stronger than (P2): for $\theta = \{\lambda_1, \lambda_2\}$ of Example 1, the graph $\mathcal{G}_{\lambda_1}^\theta$ has a switching cycle (the one presented below Figure 4), whereas \mathcal{G}_{λ_1} does not. However, it is not hard to check that (P2*) is implied by (P2) and (P3) together.

The bijection between compact non-monomial Girard proof structures and our proof structures can now be further refined: compact non-monomial Girard proof nets are in bijection with sets of linkings in our sense which satisfy (P1) and (P2*).

5 Work in progress

The equivalence relation on cut-free MLL proofs induced by their translation into cut-free MLL proof nets is canonical in sense that the equivalence corresponds to coherence in a star-autonomous category [BCST96]. We conjecture that the equivalence on cut-free MALL proofs induced by our translation into proof nets corresponds to coherence in a star-autonomous category with products (hence sums).

We are seeking a reformulation of cut that preserves the elegance of the cut-free definition, in the sense of retaining a natural translation from proofs to proof nets.

We are investigating whether the following variant of (P3) yields an alternative definition of proof net: for any switching cycle S of a set of linkings Λ , at least one $\&$ toggled in Λ is not in S .

Acknowledgements. Vaughan Pratt, for invaluable feedback during the development of this work. Paul-André Melliès, for suggesting to the first author that the search for a satisfactory notion of MALL proof net was an interesting and potentially fruitful research topic.

References

- [AJ94] S. ABRAMSKY & R. JAGADEESAN (1994): *Games and full completeness for multiplicative linear logic*. *Journal of Symbolic Logic* 59(2), pp. 543–574.
- [AM99] S. ABRAMSKY & P.-A. MELLIÈS (1999): *Concurrent games and full completeness*. In *Proc. LICS'99*, IEEE, pp. 431–442.
- [BCST96] R. F. BLUTE, J. R. B. COCKETT, R. A. G. SEELY & T. H. TRIMBLE (1996): *Natural deduction and coherence for weakly distributive categories*. *Journal of Pure and Applied Algebra* 113, pp. 229–296.
- [BS96] R. F. BLUTE & P. J. SCOTT (1996): *Linear Läuchli semantics*. *Annals of Pure & Applied Logic* 77, pp. 101–142.
- [DHPP99] H. DEVARAJAN, D. HUGHES, G. PLOTKIN & V. R. PRATT (1999): *Full completeness of the multiplicative linear logic of Chu spaces*. In *Proc. LICS'99*, IEEE, pp. 234–245.
- [DR89] V. DANOS & L. REGNIER (1989): *The structure of multiplicatives*. *Arch. Math. Logic* 28, pp. 181–203.
- [Gir87] J.-Y. GIRARD (1987): *Linear logic*. *Theoretical Computer Science* 50, pp. 1–102.
- [Gir96] J.-Y. GIRARD (1996): *Proof-nets: the parallel syntax for proof theory*. *Logic and Algebra*. Marcel Dekker.
- [HO93] J. M. E. HYLAND & C.-H. L. ONG (1993): *Fair games and full completeness for multiplicative linear logic without the MIX-Rule*. On Ong's web page.
- [Loa94] R. LOADER (1994): *Linear logic, totality and full completeness*. In *Proc. LICS'94*, IEEE, pp. 292–298.
- [Tan97] A. TAN (1997): *Full completeness for models of linear logic*. PhD thesis, University of Cambridge.

A Appendix: Separation

Throughout this appendix θ is a proof net on a cut sequent Γ .⁹ For vertices x and y of the graph \mathcal{G}_θ , write $x \rightarrow y$ iff x is an argument of y , or x is an axiom link and y is either a $\&$ on which x depends in θ or one of the two literals linked by x , thus directing the edges of \mathcal{G}_θ downwards. Property (P2) implies \mathcal{G}_θ is connected.

A subset $\Lambda \subseteq \theta$ is **saturated** if any strictly larger subset of θ toggles more $\&$'s than Λ . Clearly θ itself is saturated. For Λ a set of linkings and w a $\&$ of Γ let Λ^w denote the set of all linkings in Λ whose additive resolution does not contain the right argument of w . Write $\lambda \stackrel{w}{=} \lambda'$ if linkings $\lambda, \lambda' \in \theta$ are either equal or w is the only $\&$ toggled between $\Gamma \upharpoonright \lambda$ and $\Gamma \upharpoonright \lambda'$. It is straightforward to check that:

- (S1) If Λ is saturated and toggles w then also Λ^w is saturated.
- (S2) If Λ is saturated and toggles w and $\lambda \in \Lambda$ then $\lambda \stackrel{w}{=} \lambda_w$ for some $\lambda_w \in \Lambda^w$.
- (S3) If Λ is saturated and toggles w and $\lambda \stackrel{x}{=} \lambda'$ for $\lambda, \lambda' \in \Lambda$ then $\lambda \stackrel{w}{=} \lambda_w \stackrel{x}{=} \lambda'_w \stackrel{w}{=} \lambda'$ for some $\lambda_w, \lambda'_w \in \Lambda^w$.

LEMMA 1 *Let w be a $\&$ toggled by a saturated set $\Lambda \subseteq \theta$, and let e be an edge in \mathcal{G}_Λ originating from an axiom link a , such that $e \notin \mathcal{G}_{\Lambda^w}$. Then the jump $a \rightarrow w$ is in \mathcal{G}_Λ .*

Proof. Let e be $a \rightarrow x$. If e is not a jump, $e \notin \mathcal{G}_{\Lambda^w}$ implies $a \notin \mathcal{G}_{\Lambda^w}$. Choose $\lambda \in \Lambda$ with $a \in \lambda$. By (S2) $\lambda \stackrel{w}{=} \lambda_w$ for some $\lambda_w \in \Lambda^w$. Since $a \notin \lambda_w$ (for $a \notin \mathcal{G}_{\Lambda^w}$), the jump $a \rightarrow w$ is in \mathcal{G}_Λ .

If e is a jump, we have $\lambda, \lambda' \in \Lambda$ with $a \in \lambda$, $a \notin \lambda'$ and $\lambda \stackrel{x}{=} \lambda'$. By (S3) $\lambda \stackrel{w}{=} \lambda_w \stackrel{x}{=} \lambda'_w \stackrel{w}{=} \lambda'$ for $\lambda_w, \lambda'_w \in \Lambda^w$. Either $a \notin \lambda_w$ or $a \in \lambda'_w$, else $e \in \mathcal{G}_{\Lambda^w}$; either way, the jump $a \rightarrow w$ is in \mathcal{G}_Λ . \square

LEMMA 2 *Every non-empty union S of switching cycles in \mathcal{G}_θ has a jump out of it, i.e., for some axiom link $a \in S$ and $\&$ -vertex $w \notin S$, there is a jump $a \rightarrow w$ in \mathcal{G}_θ .*

Proof. Let Λ be a minimal saturated subset of θ with \mathcal{G}_Λ containing S . Switchings of singleton subsets of θ are cycle-free by (P2), so Λ contains at least two linkings. Let w be a $\&$ toggled by Λ that is not in any switching cycle of Λ (existing by (P3)), so $w \notin S$. Since Λ is minimal, $S \not\subseteq \mathcal{G}_{\Lambda^w}$ (using (S1)), so some edge e of S is in \mathcal{G}_Λ but not in \mathcal{G}_{Λ^w} . Without loss of generality e is an edge from an axiom link a , because for any other edge $y \rightarrow x$ in S we have $a \rightarrow z_1 \rightarrow \dots \rightarrow z_n = y \rightarrow x$ in S for some axiom link a , and $y \rightarrow x$ is in \mathcal{G}_{Λ^w} whenever $a \rightarrow z_1$ is in \mathcal{G}_{Λ^w} . By Lemma 1 the jump $a \rightarrow w$ is in \mathcal{G}_Λ , hence also in \mathcal{G}_θ . \square

⁹Readers concerned only with the cut-free case may ignore the full generality of this reference to cut, and assume that θ is a cut-free proof net on a MALL sequent Γ .

Henceforth “ $\mathfrak{Y}/\&$ ” abbreviates “ \mathfrak{Y} or $\&$ ”. A **path** from x_0 to x_n in \mathcal{G}_θ is a sequence of distinct vertices $x_0 x_1 \dots x_n$ such that for each i either $x_i \rightarrow x_{i+1}$ or $x_i \leftarrow x_{i+1}$ (note that we do not allow cycles). A **switching path** is a path in \mathcal{G}_θ that does not traverse two switch edges of any \mathfrak{Y} or $\&$ in succession. A switching path $x_0 \dots x_n$ is **strong** if it does not end by entering a $\mathfrak{Y}/\&$ along one of its switch edges (i.e., $x_{n-1} \rightarrow x_n$ only if x_n is not a $\mathfrak{Y}/\&$). Suppose paths $\pi = x_0 \dots x_n$ and $\pi' = y_0 \dots y_m$ are disjoint but for $x_n = y_0$. If π and π' are switching paths, then their composite $\pi; \pi' = x_0 \dots x_n y_1 \dots y_m$ is not necessarily a switching path (namely if $x_n = y_0$ is a $\&/\mathfrak{Y}$ and $x_{n-1} \rightarrow x_n = y_0 \leftarrow y_1$), whereas if π and π' are strong switching paths then $\pi; \pi'$ is also a strong switching path.

Write $x \Rightarrow_G y$ if the subgraph G of \mathcal{G}_θ contains a strong switching path from x to y . An **x -zone** X is a subgraph of \mathcal{G}_θ such that for all $x' \in X$, there exists y such that $x \leftarrow y \Rightarrow_X x'$.¹⁰ Given a $\mathfrak{Y}/\&$ -vertex x and a vertex y , define x **dominates** y , denoted $x \sqsupset y$, if y is in an x -zone. If x is not dominated, it is **free**.

LEMMA 3 *Domination \sqsupset is transitive.*

Proof. We show that if X is an x -zone, $y \in X$ and Y is a y -zone, then $X \cup Y$ is an x -zone. Take $z \in Y \setminus X$. We have $x \leftarrow x' \Rightarrow_X y \leftarrow y' \Rightarrow_Y z$ for some $x' \in X$ and $y' \in Y$. If the strong switching path $y' \Rightarrow_Y z$ does not intersect X , then $x' \Rightarrow_X y \leftarrow y' \Rightarrow_Y z$ constitutes a strong switching path, so we are done. Otherwise let y'' be the last vertex along $y' \Rightarrow_Y z$ that is in X . Since $y'' \in X$ we have $x \leftarrow x'' \Rightarrow_X y''$, and the sub-path of $y' \Rightarrow_Y z$ from y'' to z is a strong switching path $y'' \Rightarrow_Y z$; the composition of these paths yields $x \leftarrow x'' \Rightarrow_{X \cup Y} z$, since the only common vertex is y'' . \square

LEMMA 4 *Let C be a switching cycle in \mathcal{G}_θ containing an axiom link with a jump to a $\&$ -vertex $w \notin C$. Then w dominates every vertex of C .*

Proof. C is a w -zone. \square

LEMMA 5 *If x is in a switching cycle in \mathcal{G}_θ then $w \sqsupset x$ for some $\&$ -vertex w in no switching cycle in \mathcal{G}_θ .*

Proof. Apply Lemma 2 repeatedly, growing a collection of switching cycles one cycle at a time, until jumping to a $\&$ -vertex w that is not in a switching cycle. This must happen eventually, since (P3) implies \mathcal{G}_θ contains a $\&$ that is not in any switching cycle. The result follows by Lemma 4 and the transitivity of \sqsupset . \square

¹⁰The union of all x -zones is itself an x -zone, which could be called the **realm** of x , a concept reminiscent of the notion of **empire** of [Gir96], but different in an essential way.

LEMMA 6 *If $x \sqsupset x$ then x is in a switching cycle in \mathcal{G}_θ .*

Proof. If $x \sqsupset x$ then $x \leftarrow y \Rightarrow_X x$ for some x -zone X , hence x is in a switching cycle in \mathcal{G}_θ . \square

LEMMA 7 *Every $\mathfrak{N}/\&$ of \mathcal{G}_θ is either free or is dominated by a free $\mathfrak{N}/\&$.*

Proof. If x_0 is neither free nor dominated by a free $\mathfrak{N}/\&$ -vertex, then we can build an infinite chain $x_0 \sqsubset x_1 \sqsubset \dots$ of distinct vertices with the same property. If x_i is in a switching cycle then take x_{i+1} to be the vertex given by Lemma 5, which is not in a switching cycle; x_{i+1} is fresh otherwise $x_{i+1} \sqsubset x_{i+1}$ whence x_{i+1} is in a switching cycle (contradiction). If x_i is in no switching cycle then x_{i+1} exists since x_i is not free; x_{i+1} is fresh otherwise $x_i \sqsubset x_i$, whence x_i is in a switching cycle (contradiction). \square

COROLLARY 1 *If \mathcal{G}_θ has a $\mathfrak{N}/\&$ then it has a free $\mathfrak{N}/\&$.*

LEMMA 8 *If $x \sqsupset y_0$ and there is a path $y_0 y_1 \dots y_n$ in \mathcal{G}_θ which never enters a $\mathfrak{N}/\&$ from above (i.e., $y_{i-1} \rightarrow y_i$ only if y_i is not a $\mathfrak{N}/\&$), then $x \sqsupset y_n$.*

Proof. Let X be an x -zone containing y_0 , and let y_i be the last vertex on $y_0 y_1 \dots y_n$ that is in X . Then $x \leftarrow z \Rightarrow_X y_i$ for some z . Now $Y = X \cup \{y_i, \dots, y_n\}$ is an x -zone, since $x \leftarrow z \Rightarrow_Y y_i \Rightarrow_Y y_j$ for $j = i, \dots, n$. \square

Distinct $\mathfrak{N}/\&$ -vertices x and y of \mathcal{G}_θ are **back-to-back**, denoted $x \longleftrightarrow y$, if there is a switching path $x z_0 \dots z_n y$ in \mathcal{G}_θ such that $x \leftarrow z_0$ and $z_n \rightarrow y$, and are **face-to-face**, denoted $x \rightarrow \leftarrow y$, if there exists a path $x z_0 \dots z_n y$ in \mathcal{G}_θ such that $x \rightarrow z_0$ and $z_n \leftarrow y$, and none of the z_i are $\mathfrak{N}/\&$ -vertices (so in particular $x z_0 \dots z_n y$ is a strong switching path).

LEMMA 9 *If $x \sqsupset z$ and $y \sqsupset z$ for distinct free $\mathfrak{N}/\&$ -vertices x and y , then $x \longleftrightarrow y$.*

Proof. Let X be an x -zone containing z , so that for some x' we have $x \leftarrow x'$ and a strong switching path $\pi = x' \dots z$ in X . Let z' be the first vertex of $x\pi$ with $y \sqsupset z'$. By Lemma 8 the predecessor z'' of z' in $x\pi$ is a $\mathfrak{N}/\&$ and $z'' \leftarrow z'$, so the prefix π' of π up to z' is strong. Since $y \sqsupset z'$ there is a strong switching path $\pi'' = y_1 \dots y_n z'$ in a y -zone, with $y \leftarrow y_1$. The concatenation $x\pi'' y_n \dots y_1 y$ is a switching path (since none of the y_i is in π') witnessing $x \longleftrightarrow y$. \square

A $\mathfrak{N}/\&$ -vertex x of \mathcal{G}_θ **separates** if it is not an argument (i.e., is an outermost connective), or it is the argument of y and deleting the edge between x and y disconnects \mathcal{G}_θ .

LEMMA 10 *If a $\mathfrak{N}/\&$ -vertex x is free and does not separate, then $x \rightarrow \leftarrow y$ and $x \longleftrightarrow z$ for free y and z .*

Proof. Since x does not separate, it is in a cycle C (say clockwise) whose first (resp. last) edge is oriented out of (resp. into) x . Take y to be the first $\mathfrak{N}/\&$ reached clockwise along C from x . Then $x \rightarrow \leftarrow y$ (otherwise $y \sqsupset x$ by Lemma 8) and y is free since $y' \sqsupset y$ implies $y' \sqsupset x$ by Lemma 8, contradicting the freedom of x .

Let z be the first vertex reached anti-clockwise from x that is not dominated by x , and let z' be its predecessor. By Lemma 8, z is a $\mathfrak{N}/\&$, and $x \sqsupset z' \rightarrow z$, therefore $x \longleftrightarrow z$. If z is not free, replace it by a free $\mathfrak{N}/\&$ dominating z (hence also z') provided by Lemma 7, and appeal to Lemma 9. \square

LEMMA 11 *Let x be a $\mathfrak{N}/\&$ and let $z_0 \dots z_n$ be a switching path in \mathcal{G}_θ such that $z_0 \rightarrow x$ and $z_n \rightarrow x$. Then x dominates every vertex of $\{z_0, \dots, z_n\}$.*

Proof. $\{z_0, \dots, z_n\}$ is a x -zone. \square

SEPARATION LEMMA *If \mathcal{G}_θ has a $\mathfrak{N}/\&$ -vertex then it has a $\mathfrak{N}/\&$ -vertex that separates.*

Proof. Had \mathcal{G}_θ no separating $\mathfrak{N}/\&$ -vertex then $x_0 \rightarrow \leftarrow x_1 \longleftrightarrow x_2 \rightarrow \leftarrow x_3 \longleftrightarrow \dots$ for free $\mathfrak{N}/\&$ -vertices x_i with $x_{i+1} \neq x_i$ by Lemma 10 (x_0 exists by Corollary 1). The composition π of the paths witnessing the $\rightarrow \leftarrow$ and \longleftrightarrow relations eventually intersects itself at a vertex x , yielding a path $\pi' = x z_0 \dots z_n$ such that $\{x, z_0, \dots, z_n\}$ is a cycle. Since each witness is a switching path, π' is a switching path (by design, composition at each x_i avoids introducing consecutive switch edges of x_i). Furthermore, one of the x_i must be among the z_j . Using Lemma 5 if $\{x, z_0, \dots, z_n\}$ is a switching cycle, and Lemma 11 otherwise, this x_i is dominated, a contradiction (since x_i is free). \square

B Appendix: Proof that every cut-free proof net is the translation of a cut-free proof

With the Separation Lemma in hand, the proof that every cut-free proof net is the translation of a cut-free proof reduces to simple induction.

Let θ be a proof net on Γ . We proceed by induction on the sum the number of \mathfrak{N} 's and $\&$'s of \mathcal{G}_θ .

Base case (primary induction) Γ is $\mathfrak{N}/\&$ -free, hence θ comprises a single linking λ on Γ . We proceed by induction on the number of connectives of Γ .

- *Base case (secondary induction).* Γ contains no connectives, so $\Gamma = P_1, P_1^\perp, \dots, P_n, P_n^\perp$ for $n \geq 0$ and propositional variables P_1, \dots, P_n , and λ links the complementary literal occurrences P_i and P_i^\perp for $i = 1, \dots, n$. By (P2) $n = 1$. The axiom rule with conclusion P_1, P_1^\perp is a sequentialisation of θ .

- *Induction step (secondary induction).* Being void of \wp 's, \mathcal{G}_λ is the unique switching of λ , so by (P2) \mathcal{G}_λ is a tree.
 - Suppose $\Gamma = \Delta, A \oplus B$, with \oplus -vertex $x \in \mathcal{G}_\lambda$ corresponding to $A \oplus B$. Since $\Gamma \upharpoonright \lambda$ is an additive resolution, x is unary in \mathcal{G}_λ , *i.e.*, there is a unique $y \in \mathcal{G}_\lambda$ with $y \rightarrow x$. If this y is the left (resp. right) argument of x , consider the left (resp. right) \oplus -rule ρ with conclusion $\Delta, A \oplus B$ and hypothesis $\Gamma' = \Delta, A$ (resp. Δ, B). The linking λ on Γ also constitutes a linking λ' on Γ' , since no literals of the deleted \oplus -argument were incident with an axiom link of λ . The graph $\mathcal{G}_{\lambda'}$ is a tree, because \mathcal{G}_λ is a tree. Hence $\theta' = \{\lambda'\}$ is a proof net on Γ' . By induction, θ' is the translation of a cut-free MALL proof of Γ' , which when followed by ρ constitutes a cut-free MALL proof of Γ whose translation is θ .
 - Suppose $\Gamma = \Delta, A_0 \otimes A_1$, with \otimes -vertex $x \in \mathcal{G}_\lambda$ corresponding to $A_0 \otimes A_1$. Deleting x separates the tree \mathcal{G}_λ into a left tree T_0 and right tree T_1 whose respective conclusions define sequents Δ_0 and Δ_1 , a partitioning of Δ . Consider the \otimes -rule ρ with conclusion Γ and hypotheses Δ_0, A_0 and Δ_1, A_1 . Since \mathcal{G}_λ is a tree, no axiom link of λ goes between Δ_0, A_0 and Δ_1, A_1 , hence λ partitions to form linkings λ_0 and λ_1 , respectively, on Δ_0, A_0 and Δ_1, A_1 . Each $\theta_i = \{\lambda_i\}$ is a proof net on Δ_i, A_i since each $\mathcal{G}_{\lambda_i} = T_i$ is a tree. Appeal to the induction hypothesis with θ_0 and θ_1 , in the manner of the \oplus case above.

Induction step (primary induction) Γ has at least one \wp or $\&$. By (P2) \mathcal{G}_θ is connected.

- Suppose $\Gamma = \Delta, A \wp B$, with \wp -vertex $x \in \mathcal{G}_\theta$ corresponding to $A \wp B$. Consider the \wp -rule ρ with conclusion Γ and hypothesis $\Gamma' = \Delta, A, B$. The sequents Γ and Γ' have the same literal occurrences and essentially the same $\&$ - and additive resolutions, so θ constitutes a proof structure θ' on Γ' . The switchings of the linkings of θ' are trees, since they are obtained from those of θ by deleting x . Moreover, θ and θ' have the same subsets Λ of linkings, toggling the same $\&$'s and having the same switching cycles (because the vertex x cannot be in a switching cycle of any $\Lambda \subseteq \theta$). Therefore θ' is a proof net on Γ' . Appeal to the induction hypothesis with θ' .
- Suppose $\Gamma = \Delta, A_0 \& A_1$, with vertex $w \in \mathcal{G}_\theta$ corresponding to $A_0 \& A_1$. Consider the $\&$ -rule ρ with conclusion Γ and left and right hypotheses $\Gamma_0 = \Delta, A_0$ and $\Gamma_1 = \Delta, A_1$, respectively. Define the sets of linkings θ_i on Γ_i to comprise those linkings of θ which are

on $\Gamma_i \subseteq \Gamma$. Each θ_i is a proof net since any switching cycle of θ_i is a switching cycle of \mathcal{G}_θ . Appeal to the induction hypothesis with each θ_i .

- Suppose \mathcal{G}_θ has no \rightarrow -terminal (*i.e.* concluding) \wp or $\&$. By the Separation Lemma \mathcal{G}_θ has a $\wp/\&$ -vertex x such that the deletion of the edge $x \rightarrow y$ disconnects \mathcal{G}_θ into G_0 and G_1 .

Let G_0 be the component containing x , and let Γ_0 comprise the formula-occurrences corresponding to the \rightarrow -terminal vertices of G_0 (some formulas of Γ together with the subformula occurrence $A \& B$ corresponding to x). Define $\theta_0 = \{\lambda \upharpoonright \Gamma_0 : \lambda \in \theta\}$ on Γ_0 (each $\lambda \upharpoonright \Gamma_0$ is well-defined since no $a \in \lambda$ goes between G_0 and G_1).

Let Γ_1 be the subsequent of Γ containing the formulas corresponding to the \rightarrow -terminal vertices of G_1 . In G_1 , y is \rightarrow -initial. Form G_1^+ by adding literals P and P^\perp , the axiom link $a = \{P, P^\perp\}$, and edges $y \leftarrow P \leftarrow a \rightarrow P^\perp$. Let $\hat{\Gamma}_1$ be Γ_1 with P substituted for the subformula occurrence $A \& B$ corresponding to x , and let $\Gamma_1^+ = \hat{\Gamma}_1, P^\perp$. Define $\theta_1 = \{\lambda \upharpoonright \hat{\Gamma}_1 \cup \{a\} : \lambda \in \theta\}$ on Γ_1^+ .

Claim: $x \in \Gamma \upharpoonright \lambda$ for all $\lambda \in \theta$.

Proof. If not, there is $\lambda \in \theta$ and a $\&$ -vertex w with x in $\Gamma \upharpoonright \lambda$ but not in $\Gamma \upharpoonright \lambda_w$ for some $\lambda_w \in \theta$ such that $\lambda \stackrel{w}{=} \lambda_w$. Thus there is a jump $b \rightarrow w$ in \mathcal{G}_θ for some $b \in G_0$ with $b \in \lambda \setminus \lambda_w$. Since linkings are total on additive resolutions there exists an axiom link $c \in \lambda_w \setminus \lambda$ connecting to the formula containing x , but not satisfying $c \rightarrow \dots \rightarrow x$, so there is a jump $c \rightarrow w$ in \mathcal{G}_θ . If $w \in G_0$ then $c \rightarrow w$ is a jump from G_1 to G_0 , and if $w \in G_1$ then $b \rightarrow w$ is a jump from G_0 to G_1 ; either case violates the disconnectedness of G_0 from G_1 . ■

The claim implies that θ_0 and θ_1 are sets of linkings on Γ_0 and Γ_1^+ , respectively. Moreover, $\mathcal{G}_{\theta_0} = G_0$ and $\mathcal{G}_{\theta_1} = G_1^+$. We now check that θ_0 and θ_1 are proof nets, *i.e.*, satisfy (P1)–(P3). Since θ satisfies (P1), θ_0 (resp. θ_1) has at least one linking on every $\&$ -resolution of Γ_0 (resp. Γ_1^+). Had θ_i two distinct linkings on the same $\&$ -resolution, there would be a jump from an axiom link in G_i to a $\&$ in G_{1-i} , violating the disconnectedness of G_0 from G_1 . Thus θ_i satisfies (P1). (P2) is trivially inherited from θ . Finally, (P3) holds since any set of linkings Λ' in θ_0 or θ_1 corresponds to a set of linkings Λ in θ toggling the same $\&$'s, such that any switching cycle of Λ' is a switching cycle of Λ .

By induction θ_0 is the translation of a cut-free proof Π_0 of Γ_0 and likewise θ_1 is the translation of Π_1 . Substituting Π_0 for the axiom rule with conclusion P, P^\perp in Π_1 yields a proof whose translation is θ . □

In the case of MALL+MIX, the connectedness requirement of (P2) does not apply. This condition is used three times in the above proof. To prove that a set of linkings is the translation of a cut-free MALL+MIX proof iff it is a cut-free mix net, where a *mix net* is a set of linkings satisfying (P1)–(P3) minus the connectedness requirement of (P2), in each part of the inductive proof above, the case that \mathcal{G}_θ is not connected can be dealt with by partitioning Γ into a number of non-empty subsequents Γ_i , each harbouring a connected component of \mathcal{G}_θ . The mix net θ projects to mix nets θ_i on Γ_i , which by induction are translations of cut-free MALL+MIX proofs Π_i . By the MIX rule these combine into a proof that translates to θ .

C Appendix: Proof that eliminating a cut from a proof net yields a proof net

In this appendix we establish that cut elimination preserves (P0)–(P3). Preservation of (P0) is trivial. Preservation of (P1) for a literal or multiplicative cut is also trivial; for an additive cut it is an immediate consequence of the following lemma.

LEMMA 12 *Let $A * A^\perp$ be an additive cut in a cut sequent Γ with $A = A_0 \& A_1$ and $A^\perp = A_0^\perp \oplus A_1^\perp$ (or vice versa), and let λ, λ' be linkings of a proof net on Γ such that the cut $\&$ is the only $\&$ toggled between $\Gamma \upharpoonright \lambda$ and $\Gamma \upharpoonright \lambda'$. Then λ and λ' take the same argument of A^\perp , i.e., exactly one of A_0^\perp and A_1^\perp occurs in both $\Gamma \upharpoonright \lambda$ and $\Gamma \upharpoonright \lambda'$.*

Proof. If λ and λ' took opposite arguments of A^\perp , an axiom link above A^\perp would depend on the cut $\&$. The resulting jump yields a switching cycle of $\{\lambda, \lambda'\}$ containing the only $\&$ toggled by $\{\lambda, \lambda'\}$, in violation of (P3). \square

Preservation of (P2) is straightforward for a literal or additive cut, since switchings correspond before and after the elimination. Preservation of (P2) for a multiplicative cut is a corollary of the well-definedness of cut elimination for MLL proof nets, since the elimination of a multiplicative cut from one of our proof nets corresponds precisely to the parallel elimination of copies of the cut in the induced MLL proof nets.¹¹

¹¹To be self-contained, we give a direct proof. Let Γ (resp. Γ') be the cut sequent before (resp. after) the elimination. By definition, the linkings remain the same. We prove the following stronger result: if every switching of a linking λ on Γ is a tree, then every switching of λ on Γ' is a tree. If the eliminated cut vertex c is absent from $\lambda \upharpoonright \Gamma$, every switching of λ on Γ' is a switching of λ on Γ , hence a tree; therefore assume c is present. Let x be the eliminated \wp , with arguments x_0, x_1 , and let y be the eliminated \otimes , with arguments y_0, y_1 . Thus $x_i \rightarrow x \rightarrow c \leftarrow y \leftarrow y_i$ in Γ . Let $\hat{\Gamma}$ be the result of deleting x, c and y (and associated edges) from Γ . Claim: every switching σ of λ on $\hat{\Gamma}$ is the disjoint union of three trees, one containing the x_i , one containing y_0 , and one containing y_1 . Proof: let σ_i be the switching of λ on Γ obtained from σ by adding $x_i \rightarrow x \rightarrow c \leftarrow y$

The remainder of this appendix is devoted to the proof that cut elimination preserves (P3).

Fix a proof net θ on a cut sequent Γ . We localise the notion of domination of Appendix A from θ to any saturated set of linkings $\Lambda \subseteq \theta$. Write $x \rightarrow_\Lambda y$ if the edge $x \rightarrow y$ of \mathcal{G}_θ is in \mathcal{G}_Λ . A subgraph X of \mathcal{G}_Λ is an *x -zone under Λ* if for all $x' \in X$ there exists y with $x \leftarrow_\Lambda y \Rightarrow_X x'$; given a $\wp/\&$ -vertex $x \in \mathcal{G}_\Lambda$ and a vertex $y \in \mathcal{G}_\Lambda$, define x *dominates y in Λ* , denoted $x \sqsupset_\Lambda y$, if $y \in X$ for some x -zone X under Λ . Lemmas 2, 3, 4, 5, 6, 8, and 11 of Appendix A localise from θ to any saturated set of linkings $\Lambda \subseteq \theta$, as follows:

LOCALISED LEMMA 2 *For every non-empty union S of switching cycles in \mathcal{G}_Λ there is a jump $a \rightarrow w$ in \mathcal{G}_Λ between an axiom link $a \in S$ and a $\&$ -vertex $w \notin S$ which is toggled by Λ .*

LOCALISED LEMMA 3 *Localised domination \sqsupset_Λ is transitive.*

LOCALISED LEMMA 4 *Let C be a switching cycle in \mathcal{G}_Λ containing an axiom link with a jump to a $\&$ -vertex $w \notin C$. Then $w \sqsupset_\Lambda x$ for all vertices $x \in C$.*

LOCALISED LEMMA 5 *If x is in a switching cycle of Λ then $w \sqsupset_\Lambda x$ for some $\&$ -vertex w toggled by Λ that is in no switching cycle of Λ .*

LOCALISED LEMMA 6 *If $x \sqsupset_\Lambda x$ then x is in a switching cycle of Λ .*

LOCALISED LEMMA 8 *If $x \sqsupset_\Lambda y_0$ and there is a path $y_0 y_1 \dots y_n$ in \mathcal{G}_Λ which never enters a $\wp/\&$ from above, then $x \sqsupset_\Lambda y_n$.*

LOCALISED LEMMA 11 *Let x be a $\wp/\&$ and let $z_0 \dots z_n$ be a switching path in \mathcal{G}_Λ such that $z_0 \rightarrow_\Lambda x$ and $z_n \rightarrow_\Lambda x$. Then $x \sqsupset_\Lambda z_i$, each $0 \leq i \leq n$.*

The proof of Localised Lemma 2 is a relatively straightforward adaptation of the proof of Lemma 2; we present it in full below. The proofs of the remaining localised lemmas are obtained by making the following substitutions in the proofs of the originals in Appendix A: Λ for θ , \sqsupset_Λ for \sqsupset , \rightarrow_Λ for \rightarrow , and *zone under Λ* for *zone*.

and $y_0 \rightarrow y \leftarrow y_1$, a priori a tree; were the y_i connected by a path π in σ then $y\pi y$ would be a cycle in each σ_i ; were x_j and y_k connected by a path π in σ then $cx_j\pi y_k yc$ would be a cycle in σ_j ; were the x_i disconnected in σ then (given the disconnection of the x_i from the y_i) they would be disconnected in each σ_i . A switching of λ on Γ' is a switching of λ on $\hat{\Gamma}$ together with cuts $x_0 \rightarrow c_0 \leftarrow y_0$ and $x_1 \rightarrow c_1 \leftarrow y_1$, and is therefore (by the claim) a tree.

Proof of Localised Lemma 2. Let Λ_m be a minimal saturated subset of Λ with \mathcal{G}_{Λ_m} containing S . Switchings of singleton sets of linkings are cycle-free by (P2), so Λ_m contains at least two linkings. Let w be a $\&$ toggled by Λ_m that is not in any switching cycle of Λ_m (existing by (P3)), so $w \notin S$. Since $\Lambda_m \subseteq \Lambda$, w is certainly toggled by Λ . Since Λ_m is minimal, $S \not\subseteq \mathcal{G}_{\Lambda_m^w}$ (using (S1)), so some edge e of S is in \mathcal{G}_{Λ_m} but not in $\mathcal{G}_{\Lambda_m^w}$. Without loss of generality e is an edge from an axiom link a , because for any other edge $y \rightarrow x$ in S we have $a \rightarrow z_1 \rightarrow \dots \rightarrow z_n = y \rightarrow x$ in S for some axiom link a , and $y \rightarrow x$ is in $\mathcal{G}_{\Lambda_m^w}$ whenever $a \rightarrow z_1$ is in $\mathcal{G}_{\Lambda_m^w}$. By Lemma 1 the jump $a \rightarrow w$ is in \mathcal{G}_{Λ_m} , hence also in \mathcal{G}_{Λ} . \square

Proof that cut elimination preserves (P3). Preservation is immediate for the elimination of a literal cut $P*P^\perp$, since for every set Λ of linkings on Γ , the $\&$ -vertices toggled by Λ and the switching cycles of Λ correspond before and after the elimination. Thus consider the elimination of an additive cut $(A_0 \& A_1) * (A_0^\perp \oplus A_1^\perp)$ or a multiplicative cut $(A_0 \wp A_1) * (A_0^\perp \otimes A_1^\perp)$.

Let θ' on the cut sequent Γ' be the result of eliminating $(A_0 \& A_1) * (A_0^\perp \oplus A_1^\perp)$ or $(A_0 \wp A_1) * (A_0^\perp \otimes A_1^\perp)$ from the proof net θ on Γ . Let x be the $\&$ or \wp and y the \oplus or \otimes of the cut, let x_0, x_1 and y_0, y_1 be the arguments of x and y respectively, and let c be the cut vertex $*$ between x and y . Thus in Γ' each of c, x and y have been deleted, and cut vertices c_0 between x_0 and y_0 and c_1 between x_1 and y_1 have been added, unless one of A_0, A_0^\perp or A_1, A_1^\perp disappeared in the ‘garbage collection’ phase of additive elimination, in which case only one of c_0 or c_1 is present.

Suppose θ' fails (P3), *i.e.*, there exists a set of two or more linkings $\Lambda' \subseteq \theta'$ such that every $\&$ in Γ' toggled by Λ' is in a switching cycle of Λ' on Γ' .

LEMMA 13 *There exists a saturated set of linkings $\Lambda \subseteq \theta$ on Γ such that Λ on Γ toggles the same $\&$'s as Λ' on Γ' , except perhaps x in addition (in the case of an additive cut).*

Proof. Since cut elimination simply deletes linkings, Λ' can also be viewed as a set of linkings on Γ , and $\Lambda' \subseteq \theta$. Furthermore, Λ' on Γ toggles exactly the same $\&$'s as Λ' on Γ' , except perhaps x in addition (in the case of an additive cut). Let Λ be a minimal saturated set of linkings of θ on Γ containing Λ' . By minimality, Λ on Γ toggles the same $\&$'s as Λ' on Γ' . \square

LEMMA 14 *The vertex y is not in a switching cycle of Λ .*

Proof. If y is in a switching cycle, then by Localised Lemma 5, Λ toggles a $\&$ -vertex $w \sqsupset_\Lambda y$ in no switching cycle of Λ . We have $w \sqsupset_\Lambda x$ by Localised Lemma 8. Necessarily $w \neq x$, otherwise $w \sqsupset_\Lambda w$ and by Localised Lemma 6 w is in a switching cycle of Λ , a contradiction.

By Lemma 13, w is toggled by Λ' on Γ' , hence¹² w is in a switching cycle C of Λ' on Γ' .

Suppose C does not go through both c_0 and c_1 . Then C induces a switching cycle of Λ on Γ , still containing w , obtained by re-routing a possible passage through c_0 or c_1 to go through c instead. This yields a contradiction.

Suppose C goes through both c_0 and c_1 . Re-routing both passages to go through c instead either yields two switching cycles through c with w in one of them, a contradiction, or yields a switching cycle C_y through y and a switching path $P_x = z_0 \dots z_n$ in \mathcal{G}_Λ with $z_0 \rightarrow_\Lambda x$ and $z_n \rightarrow_\Lambda x$, such that w is either in C_y or P_x . The first possibility immediately yields a contradiction, so assume $w \in P_x$. By Localised Lemma 11, $x \sqsupset_\Lambda w$, so by transitivity (Localised Lemma 3), $w \sqsupset_\Lambda w$, hence by Localised Lemma 6, w is in a switching cycle of Λ , a contradiction. \square

LEMMA 15 *Every $\&$ -vertex $v \neq x$ toggled by Λ on Γ is in a switching cycle of Λ on Γ .*

Proof. By Lemma 13, v is toggled by Λ' on Γ' , hence¹² v is in a switching cycle C of Λ' on Γ' . Suppose C goes through c_0 and/or c_1 . By re-routing the passage(s) through c_0 and/or c_1 to go through c instead, C induces a switching cycle of Λ on Γ that contains v , in contradiction with Lemma 14. Thus C does not go through c_0 or c_1 . Hence C is also a switching cycle of Λ on Γ , containing v . \square

COROLLARY 2 *If the cut is multiplicative, every $\&$ toggled by Λ on Γ is in a switching cycle of Λ on Γ .*

Thus if the cut is multiplicative, θ fails to be a proof net, a contradiction. Henceforth we assume the cut is additive.

LEMMA 16 *The $\&$ -vertex x is the unique $\&$ toggled by Λ that is not in any switching cycle of Λ .*

Proof. Since θ is a proof net, Λ toggles a $\&$ -vertex v in no switching cycle of Λ . By Lemma 15, necessarily $v = x$. \square

There exist linkings $\lambda_1, \lambda_2 \in \Lambda'$ such that $y_0 \in \lambda_1 \upharpoonright \Gamma'$ but $y_0 \notin \lambda_2 \upharpoonright \Gamma'$, for otherwise Λ would not toggle x . Thus there is a jump $a \rightarrow_\Lambda u$ in \mathcal{G}_Λ from an axiom link a above y_0 to a $\&$ -vertex u toggled by Λ . If $u = x$, we immediately obtain a switching cycle $x, c, y, y_0, \dots, a, x$ in \mathcal{G}_Λ , a contradiction. Thus $u \neq x$, so by Lemma 15, u is in a switching cycle of Λ , and by Localised Lemma 5 and Lemma 16, $x \sqsupset_\Lambda u$. Thus x is in a switching cycle of Λ going through c, y, y_0, a and u (or a shortcut thereof, if the strong switching path from x to u intersects the path x, c, y, y_0, \dots, a), a contradiction. Hence θ' satisfies (P3).

¹²Recall that Λ' was chosen as a witness to the failure of (P3) for θ' : any $\&$ in Γ' toggled by Λ' is in a switching cycle of Λ' on Γ' .